

Patient  
 Information

# How to take your Bowel Preparation for a standard morning appointment

It is essential that your bowel is empty for this investigation. In order for your bowel to be empty please follow the instructions below:

## 7 days before your appointment

If you have **diabetes** or are taking any medication that thins your blood, other than **aspirin** (which you can remain on) please contact the relevant Endoscopy Unit on the Medication Advice Line (answer phone). The telephone number is at the end of this leaflet.

Stop taking any of the following medication:

- Imodium® (loperamide)
- Codeine phosphate
- Lomotil (co-phenotrope)
- Iron tablets
- Fybogel® (ispaghula husk)

Please note all other medications such as heart tablets and blood pressure tablets should be continued as normal, including on the day of your test.

## 2 days before your appointment

Start on a low fibre diet. Eat foods only from the following list:

- Eggs – boiled or poached
- White fish
- Chicken
- Lean meat for example beef, lamb, veal or ham
- Gravy using stock cubes (white flour or corn flour to thicken)
- White bread or rolls (no seedy bread)
- Potatoes – boiled or mashed (no skins)
- Pasta/noodles
- White rice
- Butter/margarine – use sparingly
- Clear jelly (not red) without fruit pieces

Reference No.

**GHP11136\_05\_23**

Department

**Endoscopy**

Review due

**May 2026**

## Patient Information

- Sugar/honey/syrup/treacle/jelly (e.g. bramble jelly) /jams/marmalade (without peel)
- Boiled sweets or clear mints
- Tea/coffee
- Herbal tea – clear (avoid red colours)
- Lucozade<sup>®</sup>, water, soda water
- Fizzy drinks, squashes, clear fruit juice (no bits)
- Quorn, tofu, textured vegetable protein

### Do not fry food

### Avoid seeds and sweetcorn especially

### Drink plenty of fluids

## 1 day before your appointment

At **8:00am** have a good breakfast of food taken from the low fibre list. After this do not eat **any** solid food.

Drink plenty of clear fluid (tea/ clear herbal tea/coffee/fruit squash/carbonated water drinks or plain water). You may have small amounts of milk in tea or coffee. Clear jelly (not red) is allowed.

At **2:00pm** dissolve the contents of 1 sachet of Picolax in 1 pint (500mls) of water and drink it over 10 to 20 minutes. Drink a further 2 pints (1 litre) of water.

At **6:00pm** combine and dissolve sachet A and sachet B of Moviprep in 2 pints (1 litre) of water and drink this over 1 hour.

Drink a further 2 pints (1 litre) of clear fluid before going to bed.

**Please note:** Moviprep and Picolax may be flavoured with clear fruit juices (such as apple) or clear squash (such as lemon). Moviprep can be chilled in the fridge.

**Patient  
Information**

## Morning of your appointment

At **6:00am** combine and dissolve sachet A and sachet B of Moviprep in 2 pints (1 litre) of water and drink this over 1 hour. Clear fluids only should be taken this morning – **no solid food**.

It is important that you do not become dehydrated. Clear fluids can be drunk right up to the time of your appointment.

You should expect frequent bowel actions and eventually diarrhoea.

Some cramping in your stomach is normal. Your bottom may become quite sore. Please use a barrier cream to stop any irritation. There are many products available from your local pharmacy.

## If you have any difficulties

Sometimes, the bowel preparation may cause side effects such as nausea, vomiting or abdominal pain. If you have any allergies, please check the preparation ingredients before taking.

If you feel that you will be unable to complete the bowel preparation successfully, please contact the relevant Endoscopy Unit for advice.

## Contact information

### Medication Advice Line (answer machine)

If you have any questions relating to your medication, please leave a message and a member of staff will return your call:

### Cheltenham General Hospital

Tel: 0300 422 3370

### Gloucestershire Royal Hospital

Tel: 0300 422 8232

**Patient  
Information**

**Other Endoscopy Units**

**Cirencester Hospital**

Tel: 0300 421 6284

Monday to Friday, 8:00am to 6:00pm

**Stroud General Hospital**

Tel: 0300 421 8073

Monday to Friday, 8:00am to 6:00pm

Content reviewed: May 2023